

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a *choice*.



[VIEW CATALOG](#)

AJNR

Vertigo and Hearing Loss

F.J. Wippold II and P.A. Turski

AJNR Am J Neuroradiol 2009, 30 (8) 1623-1625

<http://www.ajnr.org/content/30/8/1623>

This information is current as
of May 12, 2025.

F.J. Wippold II
P.A. Turski

Vertigo and Hearing Loss

Dizziness and Vertigo

Dizziness and vertigo (Table 1) are common clinical complaints. Vertigo is caused by a disturbed vestibular system and is subdivided into peripheral vertigo (due to failure of the end organs) or central vertigo (due to failure of the vestibular nerves or central connections to the brainstem and cerebellum).¹⁻⁵

Benign Positional Vertigo, Ménière Disease, and Peripheral Vestibular Disorders

Patients with benign positional vertigo rarely demonstrate imaging findings.^{2,4} Ménière disease manifests as paroxysmal attacks of whirling vertigo due to failure of regulation of endolymph. CT or MR imaging, or both, may be used to evaluate the vestibular aqueduct, endolymphatic duct, and sac and to rule out associated infectious or neoplastic disease.⁴⁻¹²

Vestibular neuritis and labyrinthitis may also cause vertigo. Labyrinthitis is usually viral in origin with few sequelae; however, bacterial labyrinthitis may progress to partial or complete occlusion of the lumen of the affected labyrinth, detectable on MR imaging as loss of the signal intensity of the fluid contents.^{3,4} Progressive labyrinthitis obliterans may be diagnosed on high-resolution CT.¹³ Gadolinium enhancement of the labyrinthine structures or vestibular nerves may also occur and should not be mistaken for hemorrhage.¹⁴⁻¹⁶

Superior semicircular canal dehiscence, another cause of vertigo, can be diagnosed by high-resolution coronal CT imaging of the temporal bones.¹⁷⁻¹⁹ Diseases of the internal auditory canal and cerebellopontine angle, such as tumors, are readily evaluated with CT and MR imaging techniques.

Central Vestibular Disorders

Central lesions of the brainstem or cerebellum that result in central vertigo can be readily diagnosed by MR imaging. Posterior fossa vascular disorders may be evaluated with MR angiography or conventional angiography of the posterior fossa vasculature.^{3,20,21} Cervical spondylosis, which causes vertigo by compressive osteophyte formation, may be evaluated with CT.^{3,20,21}

Sensorineural Hearing Loss

Sensorineural hearing loss (SNHL) results from the pathologic changes of inner ear structures such as the cochlea or the au-

ditary nerve¹ and is best evaluated with gadolinium-enhanced MR imaging.²²⁻²⁵

Patients with fluctuating SNHL may have congenitally enlarged vestibular aqueducts (apertures greater than 4 mm) detected by either CT or MR imaging.²⁶⁻²⁹

The imaging findings must be correlated with audiometry.^{27,28}

Initial evaluation of symmetric or unilateral SNHL requires determination of whether the site of the lesion is cochlear³⁰ or retrocochlear.³¹ Following preliminary audiometric or auditory brain response testing, patients with retrocochlear localization should have a complete MR imaging study of the head to include the internal auditory canal, temporal bones, central nuclei in the brainstem, and the auditory pathways extending upward into the cerebral hemispheres.^{22,23,32-34} Gadolinium contrast enhancement may be used. CT is sometimes diagnostic in lesions 1.5 cm or greater in diameter when dedicated techniques are used, but it does not readily detect small brainstem lesions such as infarctions or demyelination.³³⁻⁴⁰

In general, most cochlear disorders such as otosclerosis are evaluated by high-resolution CT imaging. Similarly, preoperative assessment for cochlear implants is usually best accomplished by using thin-section CT with reformatted multiplanar images. In patients with congenital etiologies for hearing loss, recent reports suggest that high-resolution MR imaging is more useful for surgical planning.^{41,42}

Conductive Hearing Loss

Conductive hearing loss results from pathologic changes of either the external or middle ear structures and is best evaluated with CT. Indications include suspected complications of acute and chronic otomastoiditis, such as cholesteatoma, and the assessment of congenital or vascular anomalies. Fistulization through the tegmen tympani of the temporal bone is usually detected by CT, though the actual involvement of the meninges and veins is better assessed by MR imaging. MR imaging is also indicated when complicated inflammatory lesions are suspected to extend into the inner ear or toward the sigmoid sinus or jugular vein. Neoplasms arising from or extending into the middle ear require the use of both techniques, as their combined data provide essential information. Vascular imaging should be performed when there is suspicion of a paraganglioma extending into the middle ear.⁴³

Trauma

CT is used extensively to delineate fractures, ossicular dislocations, fistulous communications, and facial nerve injury and to evaluate post-traumatic hearing loss.⁴⁴

This article is a summary of the complete version of this topic, which is available on the ACR Website at www.acr.org/ac. Practitioners are encouraged to refer to the complete version.

Reprinted with permission of the American College of Radiology.

Please address correspondence to Franz J Wippold II, MD, FACR, Neuroradiology Section, Mallinckrodt Institute of Radiology, 510 S Kingshighway Blvd, St. Louis, MO 63110-1076; e-mail: wippold@mir.wustl.edu; or Patrick A Turski, MD, FACR, Department of Radiology, University of Wisconsin Hospital, E1/398, 600 Highland Ave, Madison, WI 53792-0001; e-mail: pturski@uwhealth.org

Rating of Techniques: Clinical condition—vertigo and hearing loss*

	MRI Head and Internal Auditory Canal without and with Contrast	MRI Head and Internal Auditory Canal without Contrast	CT Temporal Bone without Contrast	CT Head without and with Contrast	MRA Head with or without Contrast	CTA Head
Sensorineural hearing loss, acute and intermittent vertigo	8	7	6†	3	N/A	N/A
Sensorineural hearing loss, no vertigo	8	7	5	4	N/A	N/A
Conductive hearing loss, rule out petrous bone abnormality	3	3‡	8	3	N/A	N/A
Total deafness, cochlear implant candidate, surgical planning	5	5	9	3	N/A	N/A
Fluctuating hearing loss, history of meningitis or to rule out congenital anomaly	7	7	8	4	N/A	N/A
Episodic vertigo, new onset (hours to days)	7	6	4	5	6	5
Vertigo, no hearing loss, normal findings on neurologic examination	8	7	5	4	N/A	N/A

Note:—MRI indicates MR imaging; MRA, MR angiography; CTA, CT angiography; N/A, not rated.

* Appropriateness criteria scale from 1 to 9: 1 indicates least appropriate; 9, most appropriate.

† For possible cholesteatoma with labyrinthine fistula.

‡ MR imaging is superior to CT for the detection of dural invasion and extradural extension.

Congenital and Childhood Hearing Loss

The ideal imaging method for children with unilateral or asymmetric sensory neural hearing loss is still controversial. Most reports suggest that children with unilateral or asymmetric sensory neural hearing loss should have a high-resolution temporal bone CT scan and that brain and temporal bone MR imaging be obtained in select cases. In general high-resolution CT has been shown to be efficacious for the preoperative work-up for congenital hearing loss due to aural dysplasia, congenital ossicular anomalies, large vestibular aqueduct syndrome, congenital absence of cochlear nerve, and labyrinthitis ossificans.⁴⁵⁻⁵⁴

Review Information

This guideline was originally developed in 1996. The last review and update was completed in 2008.

Appendix

Expert Panel on Neurologic Imaging: Franz J. Wippold II, MD, Co-Author and Panel Chair; Patrick A. Turski, MD, Co-Author; Rebecca S. Cornelius, MD; James A. Brunberg, MD; Patricia C. Davis, MD; Robert L. De La Paz, MD; Pr. Didier Dormont; Linda Gray, MD; John E. Jordan, MD; Suresh Kumar Mukherji, MD; David J. Seidenwurm, MD; Robert D. Zimmerman, MD; Brian Nussenbaum, MD, American Academy of Otolaryngology; Michael A. Sloan, MD, MS, American Academy of Neurology.

References

1. Bagai A, Thavendiranathan P, Detsky AS. Does this patient have hearing impairment? *JAMA* 2006;295:416–28
2. McGee SR. Dizzy patients: diagnosis and treatment. *West J Med* 1995;162:37–42
3. Phelps PD, Lloyd GA. Radiology of vertigo. In: Phelps PD, Lloyd GA, eds. *Radiology of the Ear*. St. Louis: Blackwell Scientific Publications; 1983:137–41
4. Dickens JR, Graham SS. Evaluation of the dizzy patient. *Ear Hear* 1986;7:133–37
5. Macleod D, McAuley D. Vertigo: clinical assessment and diagnosis. *Br J Hosp Med (Lond)* 2008;69:330–34
6. Albers FW, Van Weissenbruch R, Casselman JW. 3DFT-magnetic resonance imaging of the inner ear in Meniere's disease. *Acta Otolaryngol* 1994;114:595–600

7. Kraus EM, Dubois PJ. Tomography of the vestibular aqueduct in ear disease. *Arch Otolaryngol* 1979;105:91–98
8. Lorenzi MC, Bento RF, Daniel MM, et al. Magnetic resonance imaging of the temporal bone in patients with Meniere's disease. *Acta Otolaryngol* 2000;120:615–59
9. Mateijsen DJ, Van Hengel PW, Krikke AP, et al. Three-dimensional Fourier transformation constructive interference in steady state magnetic resonance imaging of the inner ear in patients with unilateral and bilateral Meniere's disease. *Otol Neurotol* 2002;23:208–13
10. Nakashima T, Naganawa S, Sugiyama M, et al. Visualization of endolymphatic hydrops in patients with Meniere's disease. *Laryngoscope* 2007;117:415–20
11. Sajjadi H, Paparella MM. Meniere's disease. *Lancet* 2008;372:406–14
12. Xenellis J, Vlahos L, Papadopoulos A, et al. Role of the new imaging modalities in the investigation of Meniere's disease. *Otolaryngol Head Neck Surg* 2000;123:114–19
13. Hasso AN, Ledington JA. Imaging modalities for the study of the temporal bone. *Otolaryngol Clin North Am* 1988;21:219–44
14. Mark AS, Seltzer S, Nelson-Drake J, et al. Labyrinthine enhancement on gadolinium-enhanced magnetic resonance imaging in sudden deafness and vertigo: correlation with audiologic and electronystagmographic studies. *Ann Otol Rhinol Laryngol* 1992;101:459–64
15. Seltzer S, Mark AS. Contrast enhancement of the labyrinth on MR scans in patients with sudden hearing loss and vertigo: evidence of labyrinthine disease. *AJNR Am J Neuroradiol* 1991;12:13–16
16. Weissman JL, Curtin HD, Hirsch BE, et al. High signal from the otic labyrinth on unenhanced magnetic resonance imaging. *AJNR Am J Neuroradiol* 1992;13:1183–87
17. Belden CJ, Weg N, Minor LB, et al. CT evaluation of bone dehiscence of the superior semicircular canal as a cause of sound- and/or pressure-induced vertigo. *Radiology* 2003;226:337–43
18. Curtin HD. Superior semicircular canal dehiscence syndrome and multi-detector row CT. *Radiology* 2003;226:312–14
19. Mong A, Loevner LA, Solomon D, et al. Sound- and pressure-induced vertigo associated with dehiscence of the roof of the superior semicircular canal. *AJNR Am J Neuroradiol* 1999;20:1973–75
20. Kikuchi S, Kaga K, Yamasoba T, et al. Slow blood flow of the vertebralbasilar system in patients with dizziness and vertigo. *Acta Otolaryngol* 1993;113:257–60
21. Norrving B, Magnusson M, Holtas S. Isolated acute vertigo in the elderly: vestibular or vascular disease? *Acta Neurol Scand* 1995;91:43–48
22. Busaba NY, Rauch SD. Significance of auditory brain stem response and gadolinium-enhanced magnetic resonance imaging for idiopathic sudden sensorineural hearing loss. *Otolaryngol Head Neck Surg* 1995;113:271–75
23. Hendrix RA, DeDio RM, Sclafani AP. The use of diagnostic testing in asymmetric sensorineural hearing loss. *Otolaryngol Head Neck Surg* 1990;103:593–98
24. Kano K, Tono T, Ushisako Y, et al. Magnetic resonance imaging in patients with sudden deafness. *Acta Otolaryngol Suppl* 1994;514:32–36
25. Weber PC, Zbar RI, Gantz BJ. Appropriateness of magnetic resonance imaging in sudden sensorineural hearing loss. *Otolaryngol Head Neck Surg* 1997;116:153–56
26. Davidson HC, Harnsberger HR, Lemmerling MM, et al. MR evaluation of ves-

- tibulocochlear anomalies associated with large endolymphatic duct and sac. *AJNR Am J Neuroradiol* 1999;20:1435–41
27. Mafee MF, Charletta D, Kumar A, et al. Large vestibular aqueduct and congenital sensorineural hearing loss. *AJNR Am J Neuroradiol* 1992;13:805–19
 28. Okumura T, Takahashi H, Honjo I, et al. Sensorineural hearing loss in patients with large vestibular aqueduct. *Laryngoscope* 1995;105:289–93, discussion 93–44
 29. Valvassori GE, Clemis JD. The large vestibular aqueduct syndrome. *Laryngoscope* 1978;88:723–28
 30. Hegarty JL, Patel S, Fischbein N, et al. The value of enhanced magnetic resonance imaging in the evaluation of endocochlear disease. *Laryngoscope* 2002;112:8–17
 31. Davidson HC. Imaging evaluation of sensorineural hearing loss. *Semin Ultrasound CT MR* 2001;22:229–49
 32. Cueva RA. Auditory brainstem response versus magnetic resonance imaging for the evaluation of asymmetric sensorineural hearing loss. *Laryngoscope* 2004;114:1686–92
 33. Selesnick SH, Jackler RK, Pitts LW. The changing clinical presentation of acoustic tumors in the MRI era. *Laryngoscope* 1993;103:431–36
 34. Gebarski SS, Tucci DL, Telian SA. The cochlear nuclear complex: MR location and abnormalities. *AJNR Am J Neuroradiol* 1993;14:1311–18
 35. Daniels RL, Swallow C, Shelton C, et al. Causes of unilateral sensorineural hearing loss screened by high-resolution fast spin echo magnetic resonance imaging: review of 1,070 consecutive cases. *Am J Otol* 2000;21:173–80
 36. Kocaoglu M, Bulakbasi N, Ucoz T, et al. Comparison of contrast-enhanced T1-weighted and 3D constructive interference in steady state images for predicting outcome after hearing-preservation surgery for vestibular schwannoma. *Neuroradiology* 2003;45:476–81
 37. Kwan TL, Tang KW, Pak KK, et al. Screening for vestibular schwannoma by magnetic resonance imaging: analysis of 1821 patients. *Hong Kong Med J* 2004;10:38–43
 38. Somers T, Casselman J, de Ceulaer G, et al. Prognostic value of magnetic resonance imaging findings in hearing preservation surgery for vestibular schwannoma. *Otol Neurotol* 2001;22:87–94
 39. Swartz JD. Lesions of the cerebellopontine angle and internal auditory canal: diagnosis and differential diagnosis. *Semin Ultrasound CT MR* 2004;25:332–52
 40. Zealley IA, Cooper RC, Clifford KM, et al. MRI screening for acoustic neuroma: a comparison of fast spin echo and contrast enhanced imaging in 1233 patients. *Br J Radiol* 2000;73:242–47
 41. Parry DA, Booth T, Roland PS. Advantages of magnetic resonance imaging over computed tomography in preoperative evaluation of pediatric cochlear implant candidates. *Otol Neurotol* 2005;26:976–82
 42. Rauch SD. Clinical practice: idiopathic sudden sensorineural hearing loss. *N Engl J Med* 2008;359:833–40
 43. Maroldi R, Farina D, Palvarini L, et al. Computed tomography and magnetic resonance imaging of pathologic conditions of the middle ear. *Eur J Radiol* 2001;40:78–93
 44. Swartz JD. Temporal bone trauma. *Semin Ultrasound CT MR* 2001;22:219–28
 45. Bamiou DE, Phelps P, Sirimanna T. Temporal bone computed tomography findings in bilateral sensorineural hearing loss. *Arch Dis Child* 2000;82:257–60
 46. Glastonbury CM, Davidson HC, Harnsberger HR, et al. Imaging findings of cochlear nerve deficiency. *AJNR Am J Neuroradiol* 2002;23:635–43
 47. Mafee MF. Congenital sensorineural hearing loss and enlarged endolymphatic sac and duct: role of magnetic resonance imaging and computed tomography. *Top Magn Reson Imaging* 2000;11:10–24
 48. McClay JE, Tandy R, Grundfast K, et al. Major and minor temporal bone abnormalities in children with and without congenital sensorineural hearing loss. *Arch Otolaryngol Head Neck Surg* 2002;128:664–71
 49. Morzaria S, Westerberg BD, Kozak FK. Evidence-based algorithm for the evaluation of a child with bilateral sensorineural hearing loss. *J Otolaryngol* 2005;34:297–303
 50. Robson CD. Congenital hearing impairment. *Pediatr Radiol* 2006;36:309–24
 51. Simons JP, Mandell DL, Arjmand EM. Computed tomography and magnetic resonance imaging in pediatric unilateral and asymmetric sensorineural hearing loss. *Arch Otolaryngol Head Neck Surg* 2006;132:186–92
 52. Tan TY, Goh JP. Imaging of congenital middle ear deafness. *Ann Acad Med Singapore* 2003;32:495–99
 53. Westerhof JP, Rademaker J, Weber BP, et al. Congenital malformations of the inner ear and the vestibulocochlear nerve in children with sensorineural hearing loss: evaluation with CT and MRI. *J Comput Assist Tomogr* 2001;25:719–26
 54. Yuen HY, Ahuja AT, Wong KT, et al. Computed tomography of common congenital lesions of the temporal bone. *Clin Radiol* 2003;58:687–93