# **Generic Contrast Agents**



Our portfolio is growing to serve you better. Now you have a *choice*.



Idiopathic Hypertrophic Spinal Pachymeningitis: Report of Two Cases with Typical MR Imaging Findings

S. Pai, C.T. Welsh, S. Patel and Z. Rumboldt

*AJNR Am J Neuroradiol* 2007, 28 (3) 590-592 http://www.ajnr.org/content/28/3/590

This information is current as of May 21, 2025.

# **CASE REPORT**

S. Pai C.T. Welsh S. Patel Z. Rumboldt

# Idiopathic Hypertrophic Spinal Pachymeningitis: Report of Two Cases with Typical MR Imaging Findings

**SUMMARY:** Two patients with a chronic progressive myelopathy were successfully surgically treated and idiopathic hypertrophic spinal pachymeningitis (IHSP) was found on histology. In both patients, an extensive extramedullary mass of low T2 signal with peripheral contrast enhancement was compressing the spinal cord on MR imaging. This imaging appearance in patients with chronic progressive myelopathy should suggest the diagnosis of IHSP.

diopathic hypertrophic spinal pachymeningitis (IHSP) is a rare inflammatory disorder that leads to spinal cord compression. We present 2 cases of this condition with a description of MR imaging findings.

# **Case Reports**

#### Case 1

A 47-year-old woman presented with a 2-year history of truncal and lower extremity weakness, numbness from the nipple line down to the feet, bowel and bladder incontinence, and increased reflexes in her lower extremities. Imaging was not performed at the initial presentation. The symptoms almost completely resolved after a year without treatment but then started progressing again.

Received February 23, 2006; accepted after revision April 12.

From the Departments of Neurosurgery (S.P., S.P.), Pathology (C.T.W.), and Radiology (Z.R.), Medical University of South Carolina, Charleston, SC.

Address correspondence to Zoran Rumboldt, MD, Medical University of South Carolina, Department of Radiology, 169 Ashley Ave, PO Box 250322, Charleston, SC 29425; e-mail: rumbolz@musc.edu.

MR imaging of the thoracic spine demonstrated an anterior extramedullary mass extending from T1 to T6 level that was of low T1 and T2 signal intensity with predominantly peripheral enhancement (Fig 1, A–C). The mass was compressing the spinal cord with resultant cord edema. Complete imaging of the neuroaxis was not performed.

After T1–T6 laminectomy, a firm tissue adherent to the ventral dura was found compressing the spinal cord, located in both intradural and extradural space. Multiple biopsy specimens were taken, but a complete resection could not be performed. A dural patch was placed posteriorly to expand the thecal sac. Chronic inflammation with fibrosis and lymphoplasmacytic infiltrate was found on histology (Fig 1*D*). There was a steady significant postoperative recovery and the patient was eventually able to ambulate with normalized bladder and bowel function.

# Case 2

A 68-year-old woman presented with a 2-month history of progressive lower and upper extremity weakness and urinary retention, bilaterally decreased sensation in lower extremities, and a poor rectal



*C*, Axial postcontrast T1-weighted image at a level similar to that of *B* reveals thick enhancement of the anterior epidural mass with central nonenhancing area (*arrowheads*). *D*, Photomicrograph shows fibrosis with plump reactive fibroblasts. Chronic inflammatory infiltrate is consisting chiefly of plasma cells (*arrows*) with additional lymphocytes (*arrowheads*) and scattered histiocytes (macrophages) (hematoxylin and eosin, original magnification  $40 \times$ ).



tone. A lower cervical laminectomy had been performed several years earlier, when the resected tissue was considered a "benign spinal tu-mor," according to the available reports.

MR imaging showed an extramedullary mass encompassing and compressing the spinal cord, hypointense on T2-weighted and isointense on T1-weighted images with attenuated, predominantly peripheral enhancement (Fig 2). Imaging of the complete neuro-axis revealed no additional abnormalities. The patient was initially treated with corticosteroids and had minimal improvement. After C6–C7 laminectomy, a large amount of intradural granular tissue attached to the dura was encountered and resected. No extradural abnormality was visible from a dorsal surgical exposure. Histology demonstrated fibrosis with chronic inflammatory reaction. The patient slowly recovered most of the neurologic deficits.

### Discussion

Idiopathic hypertrophic pachymeningitis (IHP) is a chronic progressive diffuse inflammatory fibrosis of the dura mater.<sup>1-4</sup> This rare disorder is usually found intracranially and is a diagnosis of exclusion because meningioma, lymphoma, tubercu-

Fig 2. A, Sagittal fast spin-echo (FSE) T2-weighted image shows a hypointense mass (*arrowheads*) in the anterior aspect of the spinal canal that extends from C3 to T1 level.

*B*, Corresponding sagittal postcontrast T1-weighted image with fat suppression reveals attenuated enhancement of the mass, which is predominantly peripheral with central nonenhancing portions (*arrowheads*).

losis, sarcoidosis and other diseases may present in a very similar fashion.<sup>1-5</sup> It has been proposed that Tolosa-Hunt syndrome may represent a focal manifestation of IHP.<sup>6</sup>

The spinal form is extremely rare, with few described cases, though the first report by Charcot dates back to 1869.<sup>7-13</sup> It presents as a chronic progressive disease with manifestations of radiculopathy, myelopathy, or a combination of both, attributable to the compression of neural structures or vascular compromise. It occurs alone, usually involving the cervical and thoracic dura or as a craniospinal form.<sup>1,8-13</sup> Corticosteroids have been considered the mainstay of conservative treatment; however, relapses are common, and azathioprine and cyclophosphamide have been tried with some success.<sup>1</sup> Decompression provides relief, and early surgical intervention can successfully alleviate neurologic sequelae.<sup>8-12</sup>

On MR imaging, IHSP has been described as a dural-based mass of low T2 signal intensity extending over multiple levels with strong enhancement; the linear enhancement pattern appears to show better therapeutic response than the nodular form, possibly related to less fibrosis and more vascularity.<sup>8-12</sup>

characteristics of reported cases of hubpathic hypertrophic spinal pachymeningitis (insr) that included with imaging intunings							
Study	Patient Age (years) and Sex	Spinal Levels Involved	Lesion Length in Segments	Location in Spinal Canal	Relationship to Dura	MRI Signal Intensity	MRI Enhancement Pattern
Ashkenazi et al <sup>8</sup>	65/F	T1-T5	5	Dorsal	Dural	NA	Homogenous*
Botella et al <sup>10</sup>	55/F	Cranial-C2	2	Dorsal & ventral	Dural	↓ T2 ≈T1	Peripheral
Claus et al <sup>18</sup>	31/F	T3-L1	11	Circumferential	Intradural	↓ T2 ≈T1	Homogenous
Digman et al <sup>15</sup>	70/F	All	25	Circumferential	Dural	NA	Peripheral*
Dumont et al <sup>14</sup>	30/F	C4-T3	7	Dorsal	Dural	↓ T2 ≈T1	Peripheral
Friedman and Flanders <sup>13</sup>	24/F	Cranial-C7	7	Circumferential	Dural	↓ T2 ↓ T1	Peripheral
Friedman and Flanders <sup>13</sup>	51/M	All	25	Dorsal	Dural	↓ T2 ↓ T1	Peripheral
Friedman and Flanders <sup>13</sup>	65/F	Cranial–C2	2	Circumferential	Dural	↓ T2 ↓ T1	Peripheral
Kanamori et al <sup>11</sup>	28/M	T5-L2	9	Circumferential	Dural	NA	Peripheral
Mikawa et al <sup>9</sup>	58/F	C7–T11	12	Dorsal & ventral	Dural	↓ T1	Peripheral
Park et al <sup>16</sup>	56/F	C6-T8	10	Dorsal & ventral	Dural	↓ T2 ≈T1	Peripheral
Pai et al**	47/F	T1-T6	6	Dorsal & ventral	Intra- and extradural	↓ T2 ↓ T1	Peripheral
Pai et al**	68/F	C3-T3	8	Dorsal & ventral	Intradural	↓ T2 ≈T1	Peripheral
Sridhar et al <sup>12</sup>	48/F	C7-T11	12	Circumferential	Dural	↓ T2 ↓ T1	NA
Voller et al <sup>17</sup>	54/M	Cranial-C7	7	Dorsal	Dural	NA	Peripheral

Note:—The imaging features were assessed based on inspection of the provided images or on the description of findings, in cases when the respective images were not included in the reports. ↓ indicates hypointense; ~ , isointense; \*, poor image quality; \*\*, this study; NA, not applicable (not performed or not mentioned).

Friedman and Flanders<sup>13</sup> described the peripheral enhancement that was present in all 3 patients as highly suggestive of hypertrophic pachymeningitis, which was also noted in another report.<sup>14</sup> This pattern is presumably caused by a peripheral zone of active inflammation along the lesion periphery, in contrast to attenuated fibrosis throughout the central portions.<sup>13,14</sup> In addition to fibrosis of the meninges, histology shows inflammatory cell infiltration, which is composed mainly of lymphocytes and plasma cells, especially at the surface of the dura mater.<sup>2,3</sup> We reviewed the MR images of published cases and noted that this peripheral enhancement was present in 12 of 14 patients.<sup>8-18</sup> The characteristics of reported cases are summarized in the Table. We propose that a long extramedullary mass of low T2 signal intensity with peripheral enhancement represents a specific MR imaging finding that is highly suggestive of IHSP. The specific features of the cases we presented are spontaneous temporary resolution of symptoms and disease recurrence after surgery.

#### References

- Kupersmith MJ, Martin V, Heller G, et al. Idiopathic hypertrophic pachymeningitis. Neurology 2004;62:686–94
- Riku S, Kato S. Idiopathic hypertrophic pachymeningitis. Neuropathology 2003;23:335–44
- Martin N, Masson C, Henin D, et al. Hypertrophic cranial pachymeningitis: assessment with CT and MR imaging. AJNR Am J Neuroradiol 1989;10:477–84
- Lee YC, Chueng YC, Hsu SW, et al. Idiopathic hypertrophic cranial pachymeningitis: case report with 7 years of imaging follow-up. AJNR Am J Neuroradiol 2003;24:119–23

- Parney IF, Johnson ES, Allen PBR. Idiopathic cranial hypertrophic pachymeningitis responsive to antituberculous therapy: a case report. *Neurosurgery* 1997;41:965–71
- Muthukumar N, Senthilbabu S, Usharani K. Idiopathic hypertrophic cranial pachymeningitis masquerading as Tolosa-Hunt syndrome. J Clin Neurosci 2005;12:589–92
- Charcot JM, Joffroy A. Deux cas d'atrophie musculaire progressive avec lésions de la substance grise et des faisceaux antérolatéraux de la moelle épiniére. Arch Physiol Norm Pathol 1869;2:354–67, 629–49, 744–69
- Ashkenazi E, Constantini S, Pappo O, et al. Hypertrophic spinal pachymeningitis: report of two cases and review of the literature. *Neurosur*gery 1991;28:730–32
- Mikawa Y, Watanabe R, Hino W, et al. Hypertrophic spinal pachymeningitis. Spine 1994;19:620–25
- Botella C, Orozco M, Navarro J, et al. Idiopathic chronic hypertrophic craniocervical pachymeningitis: case report. *Neurosurgery* 1994;35:1144–49
- Kanamori M, Matsui H, Terahata N, et al. Hypertrophic spinal pachymeningitis. Spine 1997;22:1787–90
- Sridhar K, Vasudevan MC. Idiopathic chronic hypertrophic pachymeningitis causing thoracic cord compression. Br J Neurosurg 2004;18:515–17
- Friedman DP, Flanders AE. Enhanced MR imaging of hypertrophic pachymeningitis. AJR Am J Roentgenol 1997;169:1425–28
- Dumont AS, Clark AW, Sevick RJ, et al. Idiopathic hypertrophic pachymeningitis: a report of two patients and review of the literature. Can J Neurol Sci 2000;27:333–34
- Digman KE, Partington CR, Graves VB. MR imaging of spinal pachymeningitis. J Comput Assist Tomogr 1990;14:988–90
- Park SH, Whang CJ, Sohn M, et al. Idiopathic hypertrophic spinal pachymeningitis: a case report. J Korean Med Sci 2001;16:683–88
- Voller B, Vass K, Wanschitz J, et al. Hypertrophic chronic pachymeningitis as a localized immune process in the craniocervical region. *Neurology* 2001;56:107–09
- Claus E, Rutgers M, Sindic C, et al. Remitting/relapsing idiopathic hypertrophic spinal pachymeningitidis: comprehensive imaging work-up and MR monitoring. Eur Radiol 2005;15:53–58